


Scheme 2

Fig. 1







Fig. 3


Recording of coordinates of particles in a bead using focal or confocal microscopy



#### 8/34


Fig. 4

Recording of coordinates of particles in a moving bead by two alternating scanning




9/34


Fig. 5a





10/34



11/34







iig. 6 B






Fig. 6








Fig. 6




Fig. 6




Fig. 6






Fig. (


 ${
m Table} \ 1$  Visual decoding of 20 beads from a library containing 400 dipeptides

	Seq	Pic	Other possible	ssible		Seq	Pic	Other possible	ossible
	CF	CF			111	HF	HL		
2	AT	AY			12	NT	NT		
3	DV	DV			13	AN	AN		
4	EV	EF			14	PH	PF		
5	FR	FD			15	PK	PK		
9	GM	GM	GF		16	}	FR	FH	VR
7	IV	IV	IP		17	NG	NG		
~	LP	LP			18	RQ	RQ	VQ	
6	MG	WW			19	NG	NG		
10	ΛM	ΛM			20	AL	AL	VL	AI

Fig. 7 A 3 Orthogonal pictures











Fig. 7 B 3 Orthogonal pictures



0,0 0,0 0.0 X1, X2 X1, X2 Y1, Y2 Y1, Y2 Z1, Z2 Z1, Z2





Fig. 8






Fig. 9





Fig. 10






Fig. 11






Fig.12




Fig. 13




Fig. 14




Fig. 15




Fig. 16




Fig. 17




Fig. 18




Fig. 19




Fig. 20




Fig. 21